



## King's Research Portal

DOI:

[10.1016/j.amjsurg.2016.09.034](https://doi.org/10.1016/j.amjsurg.2016.09.034)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Maret-Ouda, J., & Lagergren, J. (2016). The risk of mortality following secondary fundoplication in a population-based cohort study. *American Journal of Surgery*. <https://doi.org/10.1016/j.amjsurg.2016.09.034>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Official Publications of:**

- The American Surgical Congress
- The North Pacific Surgical Association
- The Association for Surgical Education
- The Association of Women Surgeons
- The Midwest Surgical Association
- The Society of Black Academic Surgeons

**Full Text:** <http://www.elsaonline.com/>  
**Submit Manuscript:** <http://www.elsevier.com/elsa>

**ELSEVIER**

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Abstract**

**Introduction:** Mortality following laparoscopic fundoplication has been found to be negligible.

However, some patients require secondary fundoplication, and the risk of mortality following such procedure is scarcely studied.

**Methods:** This nationwide Swedish population-based cohort study included all patients undergoing secondary fundoplication following primary laparoscopic fundoplication in 1997-2013, regardless of indication. Primary outcome was mortality within 90 days of surgery, and secondary outcome was postoperative length of hospital-stay.

**Results:** 9,765 patients underwent primary laparoscopic fundoplication, 540 (5.5%) patients underwent secondary fundoplication. 382 (70.7%) were conducted laparoscopically and 158 (29.3%) were conducted with an open technique. No deaths occurred within 90 days of the secondary fundoplication. Median length of stay was longer following secondary fundoplication (4.8 days, interquartile range 1.0-5.0 days), compared to primary laparoscopic fundoplication (2.5 days, interquartile range 1.0-3.0 days).

**Conclusions:** This population-based cohort study indicates that secondary fundoplication following primary laparoscopic fundoplication is a safe procedure. The longer hospital-stay following secondary fundoplication compared to primary laparoscopic fundoplication is likely explained by the higher rate of open surgical approach.

**Summary:** This Swedish population-based cohort study assessed the risk of mortality and length of stay following secondary fundoplication after primary laparoscopic fundoplication. There were no deaths within 90 days following surgery, indicating that secondary fundoplication is a safe procedure, however the length of stay was longer following secondary fundoplication compared to primary laparoscopic fundoplication.

**Keywords:** Fundoplication; Gastroesophageal reflux disease; Laparoscopic surgery; Safety

**Title:** The risk of mortality following secondary fundoplication in a population-based cohort study

**Authors:** John MARET-OUDA, M.D.<sup>1</sup>, and Jesper LAGERGREN, M.D. Ph.D.<sup>1,2</sup>

**Affiliations:** <sup>1</sup> Upper Gastrointestinal Surgery, Department of Molecular medicine and Surgery, Karolinska Institutet, and Karolinska University Hospital, Stockholm, Sweden.

<sup>2</sup> Gastrointestinal Cancer, Division of Cancer Studies, King's College London, United Kingdom.

**Corresponding Author and Requests for Reprints:** Dr. John Maret-Ouda, Upper Gastrointestinal Surgery, Department of Molecular medicine and Surgery, Karolinska Institutet, NS67, 17176 Stockholm, Sweden. E-mail: John.Maret.Ouda@ki.se, Telephone: 0046-8-51770958.

**Funding Source:** The Swedish Research Council (grant number D0547801).

**Conflict of Interest:** None of the authors has any financial or personal relationships that could inappropriately influence or bias the author's decisions, work, or manuscript.

**Category of article:** Brief Report

**Word count:** 981

## Introduction

Fundoplication is an effective and safe treatment option of severe gastroesophageal reflux disease, especially among young and physically fit patients.(1, 2) Our recently published population-based study of 8947 patients who underwent primary laparoscopic fundoplication in Sweden at the age of 18-65 years found an all-cause 90-day mortality of 0.08% (7 patients) and only 1 (0.01%) directly surgery-related death.(2) However, despite initial successful initial results, many patients get recurrent symptoms of reflux and need to undergo a secondary fundoplication. A recent assessment of 13,050 patients undergoing primary laparoscopic fundoplication in the United States, determined that 5.2% and 6.9% of the patients underwent secondary fundoplication within 5 years and 10 years, respectively.(3) Since secondary fundoplication is generally more technically demanding and more time-consuming than primary fundoplication, there is a belief that such surgery entails higher risk of severe complications, including mortality. Yet, the risk of mortality following secondary fundoplication has been scarcely studied. A recent population-based study in Denmark, including 113 patients undergoing secondary fundoplication, found a higher rate of conversion, longer hospital stay and a slightly higher 30-day postoperative mortality rate following secondary fundoplication, compared to primary fundoplication (0.81% versus 0.45%).(4) A systematic review of 20 studies, including a total of 930 patients having undergone secondary fundoplication, found only 2 perioperative deaths (0.22%).(5) The main aim of the present study was to estimate the risk of mortality within 90 days of secondary fundoplication using a population-based design, thus providing an unselected cohort.

## Materials and methods

The design of the nationwide Swedish cohort of patients aged 18-65 years who underwent primary laparoscopic fundoplication for gastroesophageal reflux diagnosis in 1997-2013 has been described elsewhere.(2) The present study was also a nationwide Swedish population-based cohort study during

the same time period, but it only included patients who had undergone secondary fundoplication (open or laparoscopic) after primary laparoscopic fundoplication, without any upper age limit and regardless of indication. The patients were identified using the Swedish Patient Registry, which records all in-hospital care in Sweden since 1987, and includes data on diagnoses, procedures, demography, and hospitalization dates.(6)

The currently used version of the Nordic Medico-Statistical Committee (NOMESCO) Classification of Surgical Procedures was introduced in the Patient Registry in 1997, which enables identification of laparoscopic and open surgical procedures separately, hence determining the start date of the study. The surgical procedures were identified using the surgical procedure codes 'JBC01' and 'JBC00', for laparoscopic and open fundoplication, respectively. The primary outcome was 90-day postoperative mortality, which is recommended in assessing postoperative mortality following upper gastrointestinal surgery.(7) Date and causes of death were retrieved from the Swedish Causes of Death Registry, which is 100% complete regarding date of death and 99.2% complete regarding main cause of death.(8) The secondary outcome was length of hospital-stay. Data from each study participant were linked using the individual 10-digit personal identity number, which is assigned to each Swedish resident upon birth or immigration, and has been validated as an effective tool for research purposes.(9) The study was approved by the Regional Ethical Review Board in Stockholm, Sweden (reference number 2014/234-31).

## Results

Among 9,765 patients who had undergone primary laparoscopic fundoplication at the age of 18 years or later, 540 patients (5.5%) underwent a secondary fundoplication with an equal number of men and women (Table 1). There were no deaths within 90 days of secondary fundoplication. Of the secondary fundoplications, 382 were laparoscopic (70.7%) and 158 were open (29.3%). The mean length of hospital

stay was longer following secondary fundoplication (4.8 days, interquartile range 1.0-5.0) compared to primary fundoplication (2.5 days, interquartile range 1.0-3.0).

## Discussion

This study found no deaths within 90 days of secondary fundoplication in 540 patients. The hospital stay following secondary fundoplication was longer compared to primary fundoplication, and nearly one third were conducted using an open surgical approach.

Among strengths of the study is the population-based design, which counteracts selection bias, the reliability of the registry data with complete coverage of surgical procedures and mortality, and the large size of the cohort. Due to the study design, based on data from national registries, the cohort reflects the outcome of all surgeries conducted in Sweden, and not limited only to specialized centers. Among weaknesses is the limited number of patients having undergone secondary fundoplication and the inability to assess the conversion rate from laparoscopic to open approach. However, the latter was not the aim of the study. Another weakness is the lack of data regarding clinical follow-up in the registers, such as quality of life or reflux symptom scores, thereby limiting conclusions regarding clinical improvement following surgery.

This study shows an even lower mortality rate than the two recent investigations examining the same topic. A population-based Danish study found a 30-day mortality rate of 0.81% (n=1) following secondary fundoplication among 124 patients.<sup>(4)</sup> A systematic review including 930 secondary fundoplications from 20 studies found 2 postoperative deaths (0.2%),<sup>(5)</sup> both deaths occurring in one study of 76 patients.<sup>(10)</sup>



There are several potential explanations for the low short-term mortality following secondary fundoplication. Recent decades have witnessed a reduction in fundoplication.<sup>(2)</sup> This reduction has led to a centralization of services to fewer surgeons with a special interest and experience in selecting suitable patients and in performing fundoplication of high quality, which should lower the risk of mortality following secondary fundoplication. In such high-volume centers, minimally invasive surgery is usually a successful approach for secondary fundoplication, and the need for open surgery is limited. Another factor that might contribute to these encouraging results is that laparoscopic fundoplication entails less scarring, fibrosis or adhesions in the operation field compared to open surgery.<sup>(11)</sup> Thus, secondary fundoplication might be less complex in an era where laparoscopy is the dominating approach for primary fundoplication. Finally, the development of postoperative care and advances in laparoscopic techniques during recent decades might contribute to explain the low mortality rates.

Secondary fundoplication has been reported to be required in approximately 5% of patients during the first five years following primary fundoplication.<sup>(3, 4)</sup> The findings of such low reoperation rates are in line with our population-based results, where the overall secondary fundoplication rate was only 5.5%. The longer hospital stay following secondary fundoplication was likely explained mainly by the fact that nearly one third had open surgery.

In conclusion, this population-based and nationwide Swedish cohort study indicates that secondary fundoplication is a safe procedure. The longer hospital stay following secondary fundoplication compared to primary fundoplication is likely explained by the higher rate of open procedures.

## References

1. Maret-Ouda J, Brusselsaers N, Lagergren J. What is the most effective treatment for severe gastro-oesophageal reflux disease? *BMJ*. 2015;350:h3169. PubMed PMID: 26108963.
2. Maret-Ouda J, Yanes M, Konings P, et al. Mortality from laparoscopic antireflux surgery in a nationwide cohort of the working-age population. *Br J Surg*. 2016 Mar 21. PubMed PMID: 26999573.
3. Zhou T, Harnsberger C, Broderick R, et al. Reoperation rates after laparoscopic fundoplication. *Surg Endosc*. 2015 Mar;29(3):510-4. PubMed PMID: 24986015.
4. Funch-Jensen P, Bendixen A, Iversen MG, Kehlet H. Complications and frequency of redo antireflux surgery in Denmark: a nationwide study, 1997-2005. *Surg Endosc*. 2008 Mar;22(3):627-30. PubMed PMID: 18071800.
5. Symons NR, Purkayastha S, Dillemans B, et al. Laparoscopic revision of failed antireflux surgery: a systematic review. *Am J Surg*. 2011 Sep;202(3):336-43. PubMed PMID: 21788005.
6. Ludvigsson JF, Andersson E, Ekbom A, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health*. 2011;11:450. PubMed PMID: 21658213. Pubmed Central PMCID: 3142234.
7. Rutegard M, Lagergren P, Johar A, Lagergren J. Time shift in early postoperative mortality after oesophagectomy for cancer. *Ann Surg Oncol*. 2015 Sep;22(9):3144-9. PubMed PMID: 25649859.
8. Johansson LA, Westerling R. Comparing Swedish hospital discharge records with death certificates: implications for mortality statistics. *Int J Epidemiol*. 2000 Jun;29(3):495-502. PubMed PMID: 10869322.
9. Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, Ekbom A. The Swedish personal identity number: possibilities and pitfalls in healthcare and medical research. *Eur J Epidemiol*. 2009;24(11):659-67. PubMed PMID: 19504049. Pubmed Central PMCID: 2773709.

10. Cowgill SM, Arnaoutakis D, Villadolid D, Rosemurgy AS. "Redo" funduplications: satisfactory symptomatic outcomes with higher cost of care. *J Surg Res.* 2007 Nov;143(1):183-8. PubMed PMID: 17950091.
11. Ten Broek RP, Bakkum EA, Laarhoven CJ, van Goor H. Epidemiology and Prevention of Postsurgical Adhesions Revisited. *Ann Surg.* 2016 Jan;263(1):12-9. PubMed PMID: 26135678.

**Table 1. Outcomes following secondary fundoplication, compared to primary fundoplication.**

	Secondary fundoplication	Primary laparoscopic fundoplication
Number of patients	540	9,765
Age, mean (IQR)	51.3 (42-60)	48.7 (39-58)
Male sex, number (%)	269 (49.8)	5,574 (57.1)
Laparoscopic technique, number (%)	382 (70.7)	9,765 (100.0)
Open technique, number (%)	158 (29.3)	N/A
Mortality rate, number (%)	0 (0.0)	N/A
Length of hospital-stay, mean (IQR)	4.8 (1.0-5.0)	2.5 (1.0-3.0)

IQR: Interquartile range, N/A: Not applicable

- This nationwide Swedish population-based cohort study included all patients undergoing secondary fundoplication following primary laparoscopic fundoplication in 1997-2013, regardless of indication.
- There were no deaths within 90 days following secondary fundoplication.
- Length of hospital stay was longer following secondary fundoplication compared to primary laparoscopic fundoplication.